




xraybinaryorbit: A Python Package for Analyzing Orbital Modulations in X-ray Binaries

Graciela Sanjurjo-Ferrín ^{1*}, Jessica Planelles Villalva ^{1*}, Jose Miguel Torrejón ^{1*}, and Jose Joaquín Rodes-Roca ^{1*}

¹ Instituto Universitario de Física Aplicada a las Ciencias y las Tecnologías, Universidad de Alicante, 03690 Alicante, Spain * These authors contributed equally.

DOI: [10.21105/joss.07220](https://doi.org/10.21105/joss.07220)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Warrick Ball](#) 

Reviewers:

- [@matteobachetti](#)
- [@renecotyfanboy](#)

Submitted: 19 June 2024

Published: 18 December 2024

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

X-ray astronomy is a young discipline, spanning no more than a few decades. The Earth's atmosphere is opaque to this type of radiation, so observations in this part of the spectrum were delayed until the space era began, when rocket launchers carrying X-ray telescopes revealed the universe from a brand-new point of view ([Giacconi et al., 1962](#)).

X-ray binaries are tremendously interesting systems. In these extreme environments, a compact object (either a neutron star, a black hole or a white dwarf) draws in matter from a companion star, producing X-ray radiation. These systems offer a unique window into extreme physics, from the effects of strong gravity and relativistic jets to the presence of intense magnetic fields ([Carroll & Ostlie, 2017](#); [Lewin et al., 1997](#); [Shapiro & Teukolsky, 1983](#)).

Orbital modulations are observed in nearly all X-ray binary systems. These variations arise from the orbital motions within the system, driven by the relative velocities of the two stars and/or their changing configurations with respect to each other and the observer.

For instance, we might detect slight oscillating shifts in the center of emission lines and/or small variations in a neutron star's spin period. This phenomenon is likely caused by the Doppler effect. Our code translates this data into orbital parameters that explain these Doppler shifts. Although the process seems simple, it becomes complex due to factors like inclination, eccentricity, periapsis, and the masses of the stars involved. If the eccentricity is greater than zero, the velocity around the orbit is not constant, making the analysis even more intricate.

When considering stellar wind (the matter accreted by compact objects), various combinations of these factors result in orbital modulations. Eccentricity causes changes in wind density throughout the orbit, affecting the amount of accreted matter. Both eccentricity and inclination influence the absorption column faced by the emitted radiation and the ionization of the stellar wind, which also varies with orbital phase.

Although these orbital modulations are conceptually simple, they are challenging to analyze. At the same time, they can provide tremendous insights into the orbital mechanics and wind properties of our systems, may help to complete the puzzle when we already have some pieces of information or to give us a list of possible scenarios compatible with our data. This is where our tools provide valuable support.

Historically, the primary limitation in this type of analysis has been the lack of resolution for detailed phase-resolved observations. However, upcoming high-resolution missions like XRISM ([XRISM Science Team, 2022](#)) and New Athena ([Barret et al., 2016](#)) promise to significantly improve the quality of these analyses. In addition to better resolution, advances in computational power have been crucial. Many of these tools have already been successfully

applied to studies using XMM-Newton and Chandra data, enabling analyses that were previously impossible (Sanjurjo-Ferrín et al., 2021, 2022, 2024).

Statement of Need

Several software packages are available for the analysis of X-ray astronomy. Some notable Python-based options include Stingray (Huppenkothen et al., 2019), which is dedicated to the timing analysis of astronomical data, and Jaxspec (Dupourqué et al., 2024), which specializes in spectral fitting using Bayesian inference. Lightkurve (Lightkurve Collaboration et al., 2018) simplifies the analysis of time-series data from space missions like Kepler and TESS. A well-known and widely used comprehensive package for general astronomy computations and data handling is Astropy (Astropy Collaboration et al., 2022).

xraybinaryorbit is a dedicated software package specifically designed for studying orbital modulations in X-ray binary systems. The tool features an intuitive interface for managing the parameters that affect orbital modulations. During the initial use, users input parameters via a form, and these are stored in a configuration file within the working directory. This file is automatically loaded in subsequent sessions, removing the need for repeated data entry (direct loads are also permitted).

The software offers two fitting methods: least squares (LS) and particle swarm optimization (PSO) Miranda (2018). The LS method is faster but may fail to converge in some cases, whereas PSO is more robust but computationally intensive. Some results obtained with this package are shown in Figures 1, 2 and 3.

Background Theory

This package primarily relies on the following theoretical frameworks:

CAK Model

The CAK model (Castor et al., 1975) describes radiation-driven winds in massive stars, with wind velocity, density, and ionization state varying with distance to the companion star.

$$\rho = \frac{\dot{M}}{4\pi v R^2}$$

Here, ρ is wind density, \dot{M} is the mass loss rate, v is the wind velocity and R is the distance to the companion. Within this package we assume a spherical, smooth and un-ionized wind distribution.

Accretion Luminosity and Ionization

Accretion powers many close binary systems (Frank et al., 2002). The accretion luminosity is given by:

$$L_{\text{ac}} = \frac{GM\dot{M}}{R}$$

where L_{ac} is the luminosity, G is the gravitational constant, M is the compact object mass, and R is the compact object radius. The ionization parameter ξ provides information about the ionization state of the stellar wind, which, in turn, affects its absorption power and the emission lines we might observe. This parameter can be approximated as:

$$\xi = \frac{L_X}{n(r_X)r_X^2}$$

Doppler Effect

The Doppler effect refers to the change in frequency or wavelength of a wave as observed by someone who is moving relative to the wave source. The Doppler velocity induced by orbital motion and the observed Doppler-shifted wavelength can be written as:

$$v_D = -r\omega \sin \phi \sin i$$

$$\lambda_D = \lambda_{\text{rest}} \left(1 + \frac{v_D}{c} \right)$$

where r is the orbital radius, ω is angular velocity, i is inclination, and λ_D and λ_{rest} are the Doppler-shifted and rest wavelengths, respectively.




Figure 1: Reconstructed trajectory of Fe XXV emitting plasma in Cen X-3, derived from the observed Doppler shifts of the emission line center.




Figure 2: Ionization map of the stellar wind in the orbital plane of 4U 0114+65, showing variations in different observed luminosities.




Figure 3: Theoretical local absorption column for a hypothetical X-ray binary system, versus both time and orbital phase, calculated through the orbit.

Acknowledgements

This research has been funded by the ASFAE/2022/02 project from the Generalitat Valenciana.

References

- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., Earl, N., Starkman, N., Bradley, L., Shupe, D. L., Patil, A. A., Corrales, L., Brasseur, C. E., Nothe, M., Donath, A., Tollerud, E., Morris, B. M., Ginsburg, A., Vaher, E., Weaver, B. A., Tocknell, J., Jamieson, W., ... Astropy Project Contributors. (2022). The Astropy Project: Sustaining and growing a community-oriented open-source project and the latest major release (v5.0) of the core package. *The Astrophysical Journal*, *935*(2), 167. <https://doi.org/10.3847/1538-4357/ac7c74>
- Barret, D., Lam Trong, T., den Herder, J.-W., & Piro. (2016). The Athena X-ray Integral Field Unit (X-IFU). In J.-W. A. den Herder, T. Takahashi, & M. Bautz (Eds.), *Space telescopes and instrumentation 2016: Ultraviolet to gamma ray* (Vol. 9905, p. 99052F). <https://doi.org/10.1117/12.2232432>
- Bonyadi, M. R., & Michalewicz, Z. (2017). Particle swarm optimization for single objective continuous space problems: A review. *Evolutionary Computation*, *25*(1), 1–54. https://doi.org/10.1162/EVCO_r_00180
- Carroll, B. W., & Ostlie, D. A. (2017). General relativity and black holes. In *An introduction to modern astrophysics* (pp. 609–652). Cambridge University Press.
- Castor, J. I., Abbott, D. C., & Klein, R. I. (1975). Radiation-driven winds in Of stars. *The Astrophysical Journal*, *195*, 157–174. <https://doi.org/10.1086/153315>
- Dupourqué, S., Barret, D., Diez, C. M., Guillot, S., & Quintin, E. (2024). jaxspec: A fast and robust Python library for X-ray spectral fitting. *Astronomy & Astrophysics*, *690*, A317. <https://doi.org/10.1051/0004-6361/202451736>
- Frank, J., King, A., & Raine, D. (2002). *Accretion power in astrophysics* (3rd ed.). Cambridge University Press.
- Giacconi, R., Gursky, H., Paolini, F. R., & Rossi, B. B. (1962). Evidence for X rays from sources outside the solar system. *Physical Review Letters*, *9*(11), 439–443. <https://doi.org/10.1103/PhysRevLett.9.439>
- Huppenkothen, D., Bachetti, M., Stevens, A. L., Migliari, S., Balm, P., Hammad, O., Khan, U. M., Mishra, H., Rashid, H., Sharma, S., Martinez Ribeiro, E., & Valles Blanco, R. (2019). Stingray: A modern Python library for spectral timing. *The Astrophysical Journal*, *881*(1), 39. <https://doi.org/10.3847/1538-4357/ab258d>
- Lewin, W. H. G., Van Paradijs, J., & Van Den Heuvel, E. P. J. (1997). *X-ray binaries* (Vol. 26). Cambridge University Press. ISBN: 9780521599344

- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., Gully-Santiago, M., Saunders, N., Cody, A. M., Barclay, T., Hall, O., Sagar, S., Turtelboom, E., Zhang, J., Tzanidakis, A., Mighell, K., Coughlin, J., Bell, K., Berta-Thompson, Z., Williams, P., Dotson, J., & Barentsen, G. (2018). *Lightkurve: Kepler and TESS time series analysis in Python*. Astrophysics Source Code Library.
- Miranda, L. J. V. (2018). PySwarms, a research-toolkit for Particle Swarm Optimization in Python. *Journal of Open Source Software*, 3. <https://doi.org/10.21105/joss.00433>
- Sanjurjo-Ferrín, G., Torrejón, J. M., Oskinova, L., Postnov, K., Rodes-Roca, J. J., Schulz, N., & Nowak, M. (2024). Revealing the state transition of Cen X-3 at high spectral resolution with Chandra. *Astronomy & Astrophysics*, 690, A360. <https://doi.org/10.1051/0004-6361/202449724>
- Sanjurjo-Ferrín, G., Torrejón, J. M., Postnov, K., Oskinova, R.-R., L., & Bernabeu, G. (2021). X-ray variability of the HMXB Cen X-3: Evidence for inhomogeneous accretion flows. *Monthly Notices of the Royal Astronomical Society*, 501(4), 5892–5909. <https://doi.org/10.1093/mnras/staa3953>
- Sanjurjo-Ferrín, G., Torrejón, J. M., & Rodes-Roca, J. J. (2022). The first X-ray spectrum of the high-mass X-ray binary XTE J1855-026 during the compact object eclipse. *Monthly Notices of the Royal Astronomical Society*, 512(1), 304–314. <https://doi.org/10.1093/mnras/stac352>
- Shapiro, S. L., & Teukolsky, S. A. (1983). *Black holes, white dwarfs and neutron stars. The physics of compact objects*. <https://doi.org/10.1002/9783527617661>
- XRISM Science Team. (2022). XRISM quick reference. *arXiv e-Prints*, arXiv:2202.05399. <https://doi.org/10.48550/arXiv.2202.05399>