




# pathways: life cycle assessment of energy transition scenarios

Romain Sacchi <sup>1</sup> and Alvaro J. Hahn-Menacho <sup>1,2</sup>

<sup>1</sup> Laboratory for Energy Systems Analysis, Centers for Energy and Environmental Sciences and Nuclear Engineering and Sciences, Paul Scherrer Institute, 5232 Villigen, Switzerland <sup>2</sup> Chair of Energy Systems Analysis, Institute of Energy and Process Engineering, ETH Zürich, 8092 Zürich, Switzerland

DOI: [10.21105/joss.07309](https://doi.org/10.21105/joss.07309)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: Frauke Wiese 

## Reviewers:

- [@kcerva](#)
- [@marc-vdm](#)

Submitted: 24 May 2024

Published: 18 November 2024

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Summary

pathways is a Python package that conducts Life Cycle Assessment (LCA) to evaluate the environmental impacts of products, sectors, or transition scenarios over time. Unlike most energy (ESM) or integrated assessment models (IAM), pathways offers a clearer view on impacts caused by a scenario by considering supply chain relations between producers and consumers, thereby addressing direct and indirect emissions. Unlike the reported emissions in ESM and IAM scenarios, which focus primarily on operation, pathways allows reporting the environmental impacts of infrastructure build-up and decommissioning. Finally, scenarios can be characterized across a wide range of indicators which are usually not included in ESM or IAM: land use, water consumption, toxicity impacts, etc.

## Statement of need

Most IAMs and ESMs project cost- or utility-optimized future scenarios within specified greenhouse gas emissions trajectories, outlining changes needed in regional energy mixes and means of transport for global warming mitigation (Riahi et al., 2017). Prospective Life Cycle Assessment (pLCA) is crucial for evaluating the environmental performance of existing and emerging production systems, with a growing body of literature in scenario-based pLCA for emerging technologies (Bisinella et al., 2021).

Extending present-day life-cycle inventories into the future using IAM outputs, initially explored by Mendoza Beltran et al. (2018) and formalized by the Python library premise (Sacchi et al., 2022), forms the methodological basis for pLCA. Efforts in pLCA focus on improving forecasting accuracy. Performing system-wide LCAs with adjusted life cycle inventories at each time step has potential to enhance sustainability assessments, broadening focus beyond greenhouse gas emissions to include broader environmental impacts like land use, water consumption, and toxicity, addressing both direct and indirect emissions. However, system-wide LCA remains challenging due to computational costs and methodological complexities, such as defining functional units based on IAM outputs and resolving double-counting issues (Vandepaer et al., 2020; Volkart et al., 2018).

Several studies characterize energy scenarios with LCA, including Gibon et al. (2015), Rauner & Budzinski (2017) and Pehl et al. (2017), who quantified ESM or IAM scenario outputs using a hybrid-LCA framework. There is also the work of Xu et al. (2020), who developed the ambitious EAFESA framework aiming for bidirectional coupling between ESM and LCA. Yet, these studies focused on specific sectors or technologies and have not yet generalized to broader scenarios and indicators, nor have they made their implementations widely available.

Beyond conventional pLCA approaches, several tools and frameworks have been developed that leverage LCA data to support further analysis, often through automation and integration

with broader modeling frameworks. For example, the ODYM-RECC framework integrates LCA data to assess resource efficiency within climate mitigation scenarios, providing insights on material demand and supply chain impacts (Pauliuk et al., 2021). Similarly, the Mat-dp tool, when supplied with suitable input data, can be used to calculate materials needed and estimate environmental impacts of transition scenarios (Cervantes Barron & Cullen, 2022, 2024). However, because these tools depend on exogenous input data, they are not designed to systematically consider the time-dependent technology mixes influencing the production system. This limits their ability to endogenously and dynamically assess evolving environmental impacts and material demand, restricting consistency with the scenario assessed.

To address these challenges, the open-source library pathways utilizes the LCA framework brightway (Mutel, 2017) to systematically evaluate environmental impacts of energy transition scenarios. pathways works with data packages containing LCA matrices adjusted to each time step of the ESM/IAM scenario, providing detailed and transparent insights into scenario environmental impacts. pathways works particularly well with data packages produced by premise, but can be used with any ESM/IAM scenarios and LCA databases. Using LCA matrices which have been modified to reflect the scenario's time-dependent technology mixes ensures a consistent and coherent characterization of the scenario.

## Description

pathways reads a data package containing scenario data, mapping information, and LCA matrices. The data package should be a zip file containing the following files:

- datapackage.json: a JSON file describing the contents of the data package
- a mapping folder containing a mapping.yaml file that describes the mapping between the IAM scenario variables and the LCA datasets
- an inventories folder containing the LCA matrices as CSV files for each time step
- a scenario\_data folder containing the scenario data as CSV files

pathways reads the scenario data files (1 in Figure 1), and iterates, for each time step and region, through technologies with a non-null production volume. For each technology, pathways retrieves the corresponding LCI dataset by looking it up in the mapping file (2 in Figure 1). The lookup indicates pathways which LCA matrices to fetch from the data package (3 in Figure 1). The LCA matrices are loaded in bw2calc (the LCA calculation module of brightway) and multiplied by the production volume (see 4 in Figure 1). The results are aggregated and saved in a dataframe, where impacts are broken down per technology, region, time step, geographical origin of impact, life-cycle stage and impact assessment method (6 in Figure 1).

Some post-processing is done on the inventory matrices, including managing double counting. Double counting occurs when resource demands are counted multiple times across interconnected system components, inflating environmental impacts. This issue is particularly relevant when the reference scenario (e.g., from an IAM) already accounts for total regional demand, such as electricity or transport. For example, if electricity and steel production are interdependent, evaluating total electricity demand as defined by the scenario may lead to overlap: electricity requires steel, and steel production, in turn, requires additional electricity beyond the initial total. This overlap results in duplicative demand estimates.

To address this, the original LCI database is adjusted by zeroing out all regional energy inputs that the energy system model accounts for and might demand during the system's life cycle, following the same workflow presented in Volkart et al. (2018) (see 5 in Figure 1). Practitioners are required to selectively cancel out overlapping activities already accounted for by the scenario. We use a modular approach in this adjustment process, where practitioners, based on their understanding of the model generating the scenario, can select specific activity categories (e.g., electricity, heat, or specific product inputs) to exclude. For instance, if the IAM models regional electricity generation, the corresponding electricity inputs in the LCA system for upstream processes are removed to prevent double counting. Returning to the electricity-steel example,

this means the practitioner would exclude electricity inputs for steel production within the LCA, as the scenario's total electricity demand already covers this requirement.

This process is implemented in the `remove_double_accounting` function, which modifies the technosphere matrix to remove redundant entries. The function identifies flagged products for removal, locates the associated rows, and zeroes out the corresponding positions taking any specified exceptions. For instance, in the electricity-steel example, the function would find the row corresponding to regional electricity and cancel out the input in the column associated with steel production, effectively preventing double counting of electricity demand. This modular approach enhances transparency and traceability, making it easier to document and track which system components are modified, ensuring consistency between the scenario outputs and the LCA.

Finally, Global Sensitivity Analysis (GSA) can be performed on the results. Currently, pathways supports the use of the SALib library for GSA (Herman & Usher, 2017; Iwanaga et al., 2022), notably the Delta Moment-Independent Measure (DMIM) method (Borgonovo, 2007), to rank the influence of the database exchanges on the results.




Figure 1: pathways workflow: from data package to impact assessment.

A detailed [example notebook](#) is available for using pathways with a sample data package.

## Impact

By systematically updating and integrating LCA matrices over time, pathways improves the accuracy and relevance of environmental impact assessments for transition scenarios. This tool fosters greater alignment between LCAs and ESM/IAM outputs, enhancing the consistency and reliability of environmental assessments across different modelling platforms.

Additionally, pathways offers a detailed and structured workflow that enables IAM modellers to incorporate LCA into their analyses. This opens new avenues for these modellers to enhance the environmental dimension of their work.

Designed to be both reproducible and transparent, pathways facilitates collaboration and

verification within the scientific community. This approach ensures that improvements in environmental impact assessments are accessible and beneficial to a broader range of stakeholders.

## Conclusion

pathways is a tool that evaluates the environmental impacts of transition scenarios over time using time-adjusted and scenario-based LCA matrices. This approach allows for characterizing the environmental impacts of a scenario across a wide range of indicators, including land use, water consumption, toxicity impacts, etc. It also allows to attribute supply chain emissions to the final energy carriers, thus providing a more detailed and transparent view of the environmental impacts of a scenario.

## Acknowledgements

The authors gratefully acknowledge the financial support from the Swiss State Secretariat for Education, Research and Innovation (SERI), under the Horizon Europe project PRISMA (grant agreement no. 101081604). The authors also thank the Swiss Federal Office of Energy (SFOE) for the support of pathways via the SWEET program for the consortium SURE.

## References

- Bisinella, V., Christensen, T. H., & Astrup, T. F. (2021). Future scenarios and life cycle assessment: systematic review and recommendations. *International Journal of Life Cycle Assessment*, 26(11), 2143–2170. <https://doi.org/10.1007/s11367-021-01954-6>
- Borgonovo, E. (2007). A new uncertainty importance measure. *Reliability Engineering & System Safety*, 92(6), 771–784. <https://doi.org/10.1016/j.res.2006.04.015>
- Cervantes Barron, K., & Cullen, J. M. (2022). Mat-dp: An open-source Python model for analysing material demand projections and their environmental implications, which result from building low-carbon systems. *Journal of Open Source Software*, 7(76), 4460. <https://doi.org/10.21105/joss.04460>
- Cervantes Barron, K., & Cullen, J. M. (2024). Using open-source tools to project bulk and critical material demand and assess implications for low-carbon energy and transport systems: Introducing Mat-dp model tools. *Resources, Conservation and Recycling*, 209, 107803. <https://doi.org/10.1016/j.resconrec.2024.107803>
- Gibon, T., Wood, R., Arvesen, A., Bergesen, J. D., Suh, S., & Hertwich, E. G. (2015). A Methodology for Integrated, Multiregional Life Cycle Assessment Scenarios under Large-Scale Technological Change. *Environmental Science and Technology*, 49(18), 11218–11226. <https://doi.org/10.1021/acs.est.5b01558>
- Herman, J., & Usher, W. (2017). SALib: An open-source python library for sensitivity analysis. *The Journal of Open Source Software*, 2(9). <https://doi.org/10.21105/joss.00097>
- Iwanaga, T., Usher, W., & Herman, J. (2022). Toward SALib 2.0: Advancing the accessibility and interpretability of global sensitivity analyses. *Socio-Environmental Systems Modelling*, 4, 18155. <https://doi.org/10.18174/sesmo.18155>
- Mendoza Beltran, A., Cox, B., Mutel, C., Vuuren, D. van, Vivanco, D. F., Deetman, S., Edelenbosch, O., Guinée, J., & Tukker, A. (2018). When the Background Matters: Using Scenarios from Integrated Assessment Models in Prospective Life Cycle Assessment. *Journal of Industrial Ecology*. <https://doi.org/10.1111/jiec.12825>
- Mutel, C. (2017). Brightway: An open source framework for Life Cycle Assessment. *The Journal of Open Source Software*, 2(12), 236. <https://doi.org/10.21105/joss.00236>

- Pauliuk, S., Fishman, T., Heeren, N., Berrill, P., Tu, Q., Wolfram, P., & Hertwich, E. G. (2021). Linking service provision to material cycles: A new framework for studying the resource efficiency–climate change (RECC) nexus. *Journal of Industrial Ecology*, 25, 260–273. <https://doi.org/10.1111/jiec.13023>
- Pehl, M., Arvesen, A., Humpenöder, F., Popp, A., Hertwich, E. G., & Luderer, G. (2017). Understanding future emissions from low-carbon power systems by integration of life-cycle assessment and integrated energy modelling. *Nature Energy*, 2(12), 939–945. <https://doi.org/10.1038/s41560-017-0032-9>
- Rauner, S., & Budzinski, M. (2017). Holistic energy system modeling combining multi-objective optimization and life cycle assessment. *Environmental Research Letters*, 12(12). <https://doi.org/10.1088/1748-9326/aa914d>
- Riahi, K., Vuuren, D. P. van, Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., KC, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., ... Tavoni, M. (2017). The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change*, 42, 153–168. <https://doi.org/10.1016/j.gloenvcha.2016.05.009>
- Sacchi, R., Terlouw, T., Siala, K., Dirnaichner, A., Bauer, C., Cox, B., Mutel, C., Daioglou, V., & Luderer, G. (2022). PRospective EnvironMental Impact asSEment (premise): A streamlined approach to producing databases for prospective life cycle assessment using integrated assessment models. *Renewable and Sustainable Energy Reviews*, 160, 112311. <https://doi.org/10.1016/j.rser.2022.112311>
- Vandepaer, L., Panos, E., Bauer, C., & Amor, B. (2020). Energy system pathways with low environmental impacts and limited costs: Minimizing climate change impacts produces environmental cobenefits and challenges in toxicity and metal depletion categories. *Environmental Science & Technology*, 54, 5081–5092. <https://doi.org/10.1021/acs.est.9b06484>
- Volkart, K., Mutel, C. L., & Panos, E. (2018). Integrating life cycle assessment and energy system modelling: Methodology and application to the world energy scenarios. *Sustainable Production and Consumption*, 16, 121–133. <https://doi.org/10.1016/j.spc.2018.07.001>
- Xu, L., Fuss, M., Poganietz, W. R., Jochem, P., Schreiber, S., Zoephel, C., & Brown, N. (2020). An Environmental Assessment Framework for Energy System Analysis (EAFESA): The method and its application to the European energy system transformation. *Journal of Cleaner Production*, 243. <https://doi.org/10.1016/j.jclepro.2019.118614>